

# DRY COAT™

## FOR USE WITH HISTICK® 2 INOCULANT FOR SOYBEANS

### Product Information

The Dry Coat™ pre-inoculation system is a component system which allows you to pre-inoculate soybean seed up to 7 days prior to planting. The system is comprised of HiStick 2 Soybean peat-based inoculant and Dry Coat liquid polymer.

### Application Rate of Dry Coat Polymer Solution:

Add 12 oz (355 ml) of Dry Coat Polymer concentrate to 108 oz (3.2L) of clean, non-chlorinated water to prepare a total volume of 120 fl. oz. (3.55L). Apply diluted polymer at the minimum rate of 6 oz/100 lbs (177 ml/45 kg) of seed to treat a total of 2,000 lbs (907 kg).

### Application Steps For Dry Coat Polymer Pre-Inoculant System With HiStick 2 Inoculant

1. Application of Dry Coat Polymer Solution  
Calibrate pump to deliver 120 oz (3.55L) of polymer solution per 2000 lbs (907 kg) of soybean seed (6 oz/100 lbs [177 ml/45 kg]). Ensure even coverage of the soybean by matching seed flow rate with dosage rate of polymer solution.
2. Apply HiStick 2 Inoculant at the rate of 56 oz per 2000 lbs (1.6 kg/907 kg) of soybean seed immediately following seed coverage with Dry Coat Polymer solution. This should be accomplished by metering the HiStick 2 via EnviroPac, Gandy or other dry metering device to ensure even coverage to the soybeans.
3. Allow Seeds to Air Dry Prior to Bagging/Bulk Handling. Plant soybeans within 7 days of application to maintain viable rhizobia for effective nodulation.

### Seed Treatment Compatibility

Dry Coat treated soybeans are compatible with ApronMaxx® RTA™ and Bean Pak and must be planted within 7 days of application to maintain viable rhizobia numbers for effective nodulation.

### Storage/Precautions

Store Dry Coat Polymer Concentrate between 40-77°F (5-25°C) in dry conditions. Store HiStick 2 Inoculant as directed on label. Treated seeds should be stored in a cool shaded area (77°F/25°C or less) under cover. Do not allow direct exposure to sunlight for extended periods of time.

ApronMaxx® is a registered trademark and RTA™ is a trademark of Syngenta Group Company  
HiStick® is a registered trademark and DryCoat™ is a trademark of Becker Underwood Inc.

Revised 1/24/03

801 Dayton Avenue • Ames, Iowa 50010 USA • 800-232-5907 • 515-232-5907 • fax 515-232-5961

www.beckerunderwood.com

